
Atlantic Coast
Environmental
Indicators
Consortium

Emerging Indicators of Human and Climatically- Induced change in Coastal Ecosystems Hans Paerl, UNC-CH Institute of Marine Sciences Morehead City, NC www.marine.unc.edu/Paerllab

Plum Island LTER
Chesapeake Bay
Neuse/Pamlico
North Inlet

www.aceinc.org

With lots of help from:
Nathan Hall
Larry Harding
Ed Houde
Alan Joyner
Dave Kimmel
Rick Luettich
Ben Peierls
Jay Pinckney
Karen Rossignol
Lexia Valdes-Weaver
& many Others






Regional & national coastal indicators for assessing ecosystem responses to anthropogenic and naturally-induced change

What should they do?

- •Detect, quantify and predict ecosystem change for a range of estuaries varying in trophic state and water residence time
 - Clarify processes underlying change
- Evaluate human vs. natural impacts on WQ and habitat conditions
 - •Complement environmental monitoring and assessment tools

Examples (only enough time for some bio-indicators)

- Plant/productivity-based indicators
 - Trophodynamic indicators
 - Remote sensing for "scaling up"
- Be available and useful to WQ/habitat management agencies







Example: Neuse River EstuaryPamlico Sound Excessive N loading → eutrophication → hypoxia → WQ/habitat decline









N loading and algal production (Chl a) in the Neuse R. Estuary



The connection to Hypoxia



Hypoxia and Fish kills in the Neuse River Estuary 1994-2001 Data Sources: ModMon Project & NC DENR-DWQ




Fish kill data base: http://www.esb.enr.state.nc.us:80/Fishkill/fishkillmain.htm

Need: Reduce Estuarine Primary Production (Chl a) by Establishing an N Input Threshold


Scientific Consensus/Recommendation: 30% N Input Reduction (based on 1990-1995 loads)

Indicator: Using photopigments to assess algal growth response to N reductions (i.e. mandated 30% N input reduction = TMDL)

Asking the Phytoplankton: Dilution Bioassays



Seasonal Effect of 30% Reduction in N Concentration 84 Hour Incubation



Piehler et al. 2001